

Product Data Sheet

mmu-miR-185-5p miRNA Antagomir

| Catalog # | Source | Reactivity | Applications |
|-----------------------|-----------|---------------------|--|
| CIN0052 | Synthetic | Μ | |
| Description | Syn | thetic miRNA Ant | agomir is used to inhibit the activity of target mmu-miR-185-5p |
| | mR | NA. | |
| Specificity | Ant | agomir is chemic | ally-modified single-strand miRNA inhibitor functioning by |
| | blo | cking miRNA regu | lation of target gene expression efficiently. They are synthesized |
| | to r | educe the ability | of endogenous miRNAs to silence target mRNA transcripts. |
| | The | y can down-regu | late the corresponding endogenous miRNAs. Our miRNA |
| | ant | agomir is single-s | trand miRNA inhibitor carrying the chemically modifications |
| | fun | ctioning by blocki | ing miRNA regulation of target gene expression efficiently. |
| Form | Lyo | philized powder | |
| Gene Symbol | mn | nu-miR-185-5p | |
| Accession No. | MI | MAT0000214 | |
| Components | Thi | synthetic miRNA | A is based on the mature miRNA sequence. The strand of the |
| | ant | agomir has 2 pho | sphorothioates at the 5' end, 4 phosphorothioates, 1 |
| | chc | lesterol group at | the 3' end, and full-length nucleotide 2'-methoxy modification. |
| | Sta | oility of miRNA ar | ntagomir appears to be significantly higher than miRNA |
| | inh | ibitors. It exhibi | ts enhanced cellular uptake, stability and regulatory activity and |
| | is re | ecommended for | miRNA functional studies in vitro and in vivo. |
| Directions for | Use Brie | efly centrifuge tub | pes containing miRNA antagomir to ensure that the miRNA |
| | pel | et is located at th | ne bottom of the tube. Dissolve miRNA antagomir to a |
| | con | venient stock cor | ncentration using the recommended volume of DEPC H2O (or |
| | RNa | ase-free water). F | or example: dissolve 10 nmol miRNA antagomir to 20 μM using |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com