

# **Product Data Sheet**

# Anti-GADD153 (Phospho-S30) Antibody

Catalog # Source Reactivity Applications

CPA1335 Rabbit H, M, R, Mk, P WB, IH, IF/IC

**Description** Rabbit polyclonal antibody to GADD153 (Phospho-S30)

Immunogen KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding

S30 of human GADD153 protein. The exact sequence is proprietary.

**Purification** The antibody was purified by immunogen affinity chromatography.

**Specificity** Recognizes endogenous levels of GADD153 protein only when phosphorylated at

S30.

**Clonality** Polyclonal

**Conjugation** 

Form Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,

and 0.01% sodium azide.

Dilution WB (1/500 - 1/1000), IH (1/50 - 1/100), IF/IC (1/50 - 1/200)

Gene Symbol DDIT3

Alternative Names CHOP; CHOP10; GADD153; DNA damage-inducible transcript 3 protein; DDIT-3;

C/EBP zeta; C/EBP-homologous protein; CHOP; C/EBP-homologous protein 10;

CHOP-10; CCAAT/enhancer-binding protein homologous protein; Growth arrest and

DNA damage-inducible protein GADD153

Entrez Gene 1649 (Human); 13198 (Mouse)

SwissProt P35638 (Human); P35639 (Mouse); Q62857 (Rat)

Storage/Stability Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid

freeze/thaw cycles.


Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**


WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com




# **Product Data Sheet**



Western blot analysis of GADD153 (Phospho-S30) expression in HeLa (A), Raw264.7 (B), rat spleen (C) whole cell lysates. (Predicted band size: 19 kD; Observed band size: 27 kD)



Immunohistochemical analysis of GADD153 (Phospho-S30) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of GADD153 (Phospho-S30) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**