

Product Data Sheet

LRBA siRNA (Human)

Catalog #	Source	Reactivity	Applications			
CRH0702	Synthetic	н	RNAi			
Description	siRNA	to inhibit LRBA expre	ession using RNA interference			
Specificity	LRBA	LRBA siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expressio	on.			
Form	Lyoph	ilized powder				
Gene Symbol	LRBA	LRBA				
Alternative N	ames BGL; (BGL; CDC4L; LBA; Lipopolysaccharide-responsive and beige-like anchor protein;				
	Beige-	like protein; CDC4-lik	e protein			
Entrez Gene	987 (H	luman)				
SwissProt	P5085	P50851 (Human)				
Purity	> 97%	> 97%				
Quality Contr	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid				
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.				
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	human LRBA gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can				
	be tra	be transfected individually or pooled together to achieve knockdown of the target				
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	LRBA	siRNA (Human) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

LRBA siRNA (Human) - B	5 nmol x 1	5 nmol x 2
LRBA siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com