

Product Data Sheet

DDIT3 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|----------------|-----------|---|---|--|--|
| CRH1162 | Synthetic | н | RNAi | | |
| Description | siRNA | A to inhibit DDIT3 exp | ression using RNA interference | | |
| Specificity | DDIT | 3 siRNA (Human) is a | target-specific 19-23 nt siRNA oligo duplexes designed to | | |
| | knocl | k down gene expressi | on. | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | DDIT | 3 | | | |
| Alternative Na | ames CHOF | CHOP; CHOP10; GADD153; DNA damage-inducible transcript 3 protein; DDIT-3; | | | |
| | C/EBI | P zeta; C/EBP-homolo | gous protein; CHOP; C/EBP-homologous protein 10; | | |
| | CHOF | P-10; CCAAT/enhance | r-binding protein homologous protein; Growth arrest and | | |
| | DNA | damage-inducible pro | otein G | | |
| Entrez Gene | 1649 | (Human) | | | |
| SwissProt | P356 | 38 (Human) | | | |
| Purity | > 97% | 6 | | | |
| Quality Contro | ol Oligo | nucleotide synthesis | is monitored base by base through trityl analysis to ensure | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently purified by affinity-solid | | |
| | phase | e extraction. The anne | ealed RNA duplex is further analyzed by mass | | |
| | spect | rometry to verify the | exact composition of the duplex. Each lot is compared to | | |
| | the p | revious lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | |
| Components | We o | ffers pre-designed se | ts of 3 different target-specific siRNA oligo duplexes of | | |
| | huma | an DDIT3 gene. Each v | ial contains 5 nmol of lyophilized siRNA. The duplexes can | | |
| | be tra | ansfected individually | or pooled together to achieve knockdown of the target | | |
| | gene, | , which is most comm | only assessed by qPCR or western blot. | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|-------------------------|--------------|--------------|
| DDIT3 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
| DDIT3 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| DDIT3 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 μl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| | | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com