

Product Data Sheet

DPH1 siRNA (Human)

| Source | Reactivity | Applications | | |
|-----------|---|---|--|--|
| Synthetic | н | RNAi | | |
| siRNA | to inhibit DPH1 expr | ession using RNA interference | | |
| DPH1 | DPH1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| knock | down gene expressio | on. | | |
| Lyoph | nilized powder | | | |
| DPH1 | DPH1 | | | |
| mes DPH2 | DPH2L; DPH2L1; OVCA1; Diphthamide biosynthesis protein 1; DPH1 homolog; | | | |
| HsDp | h1; Diphthamide bios | ynthesis protein 2 homolog-like 1; DPH-like 1; DPH2-like | | |
| 1; Dip | ohthamide biosynthes | is protein 2-like; Ovarian cancer-associated gene 1 | | |
| prote | in | | | |
| 1801 | 1801 (Human) | | | |
| Q9BZ | Q9BZG8 (Human) | | | |
| > 97% | 6 | | | |
| ol Oligo | nucleotide synthesis i | s monitored base by base through trityl analysis to ensure | | |
| appro | opriate coupling efficie | ency. The oligo is subsequently purified by affinity-solid | | |
| phase | e extraction. The anne | ealed RNA duplex is further analyzed by mass | | |
| spect | rometry to verify the | exact composition of the duplex. Each lot is compared to | | |
| the p | revious lot by mass sp | ectrometry to ensure maximum lot-to-lot consistency. | | |
| We o | ffers pre-designed set | s of 3 different target-specific siRNA oligo duplexes of | | |
| huma | an DPH1 gene. Each vi | al contains 5 nmol of lyophilized siRNA. The duplexes can | | |
| be tra | ansfected individually | or pooled together to achieve knockdown of the target | | |
| gene, | which is most comm | only assessed by qPCR or western blot. | | |
| | Synthetic siRNA DPH1 knock Lyoph DPH1 DPH1 DPH1 DPH1 1; Dip prote 1801 Q9BZ > 97% Oligo appro phase spect the p We o huma | Synthetic H siRNA to inhibit DPH1 expr DPH1 siRNA (Human) is a to knock down gene expression Lyophilized powder DPH1 Imes DPH2L; DPH2L1; OVCA1; D HsDph1; Diphthamide biosynthesis protein 1801 (Human) Q9BZG8 (Human) > 97% Oligonucleotide synthesis i appropriate coupling efficies phase extraction. The annes Spectrometry to verify the the previous lot by mass sp We offers pre-designed set human DPH1 gene. Each vi be transfected individually | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|------------------------|--------------|--------------|
| DPH1 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
| DPH1 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| DPH1 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| | | 50 nM | 5 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com