

Product Data Sheet

ERG siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRH1451	Synthetic	Н	RNAi		
Description siRNA to inhibit ERG expression using RNA interference					
Specificity	ERG s	ERG siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knocl	k down gene expressio	on.		
Form	Lyopł	hilized powder			
Gene Symbo	ERG	ERG			
Alternative Names Transcriptional regulator ERG			RG; Transforming protein ERG		
Entrez Gene 2078 (Human)					
SwissProt	P113	P11308 (Human)			
Purity > 97%					
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			trityl analysis to ensure		
	appro	opriate coupling efficion	ency. The oligo is subsequently puri	fied by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass sp	pectrometry to ensure maximum lo	t-to-lot consistency.	
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human ERG gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can			
	be tra	ansfected individually	or pooled together to achieve know	ckdown of the target	
gene, which is most commonly assessed by qPCR or wester			only assessed by qPCR or western b	olot.	
Component		15 nmol	30 nmol		
	ERG	siRNA (Human) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

ERG siRNA (Human) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DE	PC Water	1 ml x 1	1 ml x 2
Ne	egative Control	2.5 nmol x 1	2.5 nmol x 2
ER	G siRNA (Human) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com