

# **Product Data Sheet**

## **GNGT2 siRNA (Human)**

| Catalog #     | Source  | Reactivity  | Applications                        |                             |  |
|---------------|---|---|-------------------------------------|-----------------------------|--|
| CRH1864       | Synthetic   | н   | RNAi                                |                             |  |
| Description   | siRNA   | to inhibit GNGT2 ex   | pression using RNA interference     |                             |  |
| Specificity   | GNGT  | GNGT2 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to  |                                     |                             |  |
|               | knock   | down gene expressi  | on.                                 |                             |  |
| Form          | Lyoph   | ilized powder   |                                     |                             |  |
| Gene Symbol   | GNGT  | GNGT2   |                                     |                             |  |
| Alternative N | ames GNG8   | GNG8; GNG9; GNGT8; Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit        |                                     |                             |  |
|               | gamn  | gamma-T2; G gamma-C; G-gamma-8; G-gamma-9; Guanine nucleotide binding               |                                     |                             |  |
|               | prote   | in gamma transducin   | g activity polypeptide 2            |                             |  |
| Entrez Gene   | 2793  | 2793 (Human)  |                                     |                             |  |
| SwissProt     | 0146  | O14610 (Human)  |                                     |                             |  |
| Purity        | > 97%   | > 97%   |                                     |                             |  |
| Quality Contr | Oligonucleotide synthesis is monitored base by base through trityl analys |   |                                     | h trityl analysis to ensure |  |
|               | appro   | priate coupling effici  | ency. The oligo is subsequently pu  | urified by affinity-solid   |  |
|               | phase   | e extraction. The ann   | ealed RNA duplex is further analyz  | zed by mass                 |  |
|               | spect   | spectrometry to verify the exact composition of the duplex. Each lot is compared to |                                     |                             |  |
|               | the p   | revious lot by mass s   | pectrometry to ensure maximum       | lot-to-lot consistency.     |  |
| Components    | We of   | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of  |                                     |                             |  |
|               | huma  | n GNGT2 gene. Each  | vial contains 5 nmol of lyophilized | d siRNA. The duplexes       |  |
|               | can b   | e transfected individ   | ually or pooled together to achiev  | e knockdown of the          |  |
|               | targe   | target gene, which is most commonly assessed by qPCR or western blot.               |                                     |                             |  |
|               | Com   | ponent  | 15 nmol                             | 30 nmol                     |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



# **Product Data Sheet**

| GNGT2 siRNA (Human) - A | 5 nmol x 1   | 5 nmol x 2   |
|-------------------------|--------------|--------------|
| GNGT2 siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
| GNGT2 siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control        | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water              | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μΙ         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μl         |
|         |              | 10 nM               | 0.25 μl       | 1 μΙ         |
|         |              | 100 nM              | 5 μl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
|         |              | 100 nM              | 10 µl         | 5 μΙ         |
| 6-well  | 2 ml         | 50 nM               | 5 μl          | 5 μl         |
|         |              | 10 nM               | 1 μΙ          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

## **Product Data Sheet**

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com