

Product Data Sheet

HOXA13 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRH2196	Synthetic	н	RNAi		
Description siRNA to i		to inhibit HOXA13 ex	inhibit HOXA13 expression using RNA interference		
Specificity	HOXA	HOXA13 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expres	sion.		
Form	Lyoph	ilized powder			
Gene Symbol	HOXA	HOXA13			
Alternative N	ames HOX1.	HOX1J; Homeobox protein Hox-A13; Homeobox protein Hox-1J			
Entrez Gene	3209 (3209 (Human)			
SwissProt	P3127	P31271 (Human)			
Purity > 97%					
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis t			gh trityl analysis to ensure		
	appro	priate coupling efficie	ency. The oligo is subsequently p	ourified by affinity-solid	
	phase	extraction. The anne	aled RNA duplex is further analy	yzed by mass	
	spectr	ometry to verify the	exact composition of the duplex	. Each lot is compared to	
	the pr	evious lot by mass sp	ectrometry to ensure maximum	n lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	n HOXA13 gene. Each	vial contains 5 nmol of lyophiliz	zed siRNA. The duplexes	
	can be	e transfected individu	ally or pooled together to achie	ve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	HOXA	A13 siRNA (Human) -	A 5 nmol x 1	5 nmol x 2	

HOXA13 siRNA (Human) - B 5 nmol x 1 5 nmol x 2 Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-

Application Rey: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunohistochemistry,

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DEPC Water	1 ml x 1	1 ml x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
HOXA13 siRNA (Human) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com