

Product Data Sheet

Goat Anti-Human IgG gamma chain - FITC

Catalog #	Source	e Reactivity	Applications		
CSA2305	Goat	Н	E, IH, IF, FC		
Description		Goat Polyclonal Secondary Antibody to Human IgG gamma chain FITC			
Immunogen		Human IgG gamma chain			
Purification		The antibody was isolated f	rom antisera by immunoaffinity chromatography using		
		antigens coupled to agarose	beads.		
Specificity		By immunoelectrophoresis and ELISA this antibody reacts specifically with Human			
		lgG gamma chain. No antibo	ody was detected against non immunoglobulin serum		
		proteins.			
Clonality		Polyclonal			
Conjugation		FITC			
Form		1 mg/ml. Liquid in 0.01M Pl	nosphate Buffered Saline, pH 7.2, containing 1% BSA,		
		50% glycerol, 0.05% Sodium	Azide		
Dilution		IH (1/100 - 1/1000), IF (1/100	- 1/1000), FC (1/1000 - 1/4000), E (Use at an assay		
		dependent concentration)			
Fluorophore		Ex/Em = 493⁄528 nm			
Storage/Stabi	lity	Shipped at 4°C. Upon delive	ry aliquot and store at -20°C for one year. Avoid		
		freeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com