

Product Data Sheet

Mouse Anti-Dog IgM mu chain - TRITC

Catalog #	Source	Reactivity	Applications		
CSA2672	Mouse	D	E, IH, IF, FC		
Description	Ν	Mouse Monoclonal Secondary Antibody to Dog IgM mu chain TRITC labeled			
Immunogen	C	Oog IgM mu chain			
Purification	Т	The antibody was isolated from ascitic by immunoaffinity chromatography using			
	а	intigens coupled to agaros	e beads.		
Specificity	B	By immunoelectrophoresis	and ELISA this antibody reacts specifically with Dog IgM		
	n	nu chain. No antibody wa	s detected against non immunoglobulin serum proteins.		
Clonality	Ν	Monoclonal			
Conjugation	Т	RITC			
Form	1	. mg/ml. Liquid in 0.01M I	Phosphate Buffered Saline, pH 7.2, containing 1% BSA,		
	5	0% glycerol, 0.02% Sodiu	n Azide		
Dilution	II	H (1/100 - 1/1000), IF (1/10	0 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay		
	d	lependent concentration)			
Fluorophore	E	x/Em = 595/613 nm			
Storage/Stabi	lity S	hipped at 4 $^\circ$ C. Upon del	ivery aliquot and store at -20 $^\circ$ C for one year. Avoid		
	f	reeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com