

Anti-CD62E/CD62P Antibody-Biotin labeled

Catalog #	Source	Reactivity	Applications
CFA8971	Mouse	H	IF, FC
Description	Mouse monoclonal antibody Biotin labeled to CD62E/CD62P		
Immunogen	TNF stimulated human endothelial cells		
Purification	The antibody was purified by affinity chromatography.		
Specificity	Recognizes human CD62E/CD62P		
Clonality	Monoclonal (clone: 1.2B6)		
Conjugation	Biotin		
Form	Mouse IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.		
Dilution	10 µl / assay		
Gene Symbol	SELE; SELP		
Alternative Names	ELAM1; E-selectin; CD62 antigen-like family member E; Endothelial leukocyte adhesion molecule 1; ELAM-1; Leukocyte-endothelial cell adhesion molecule 2; LECAM2; CD62E; GMRP; GRMP; P-selectin; CD62 antigen-like family member P; Granule membrane protein 140		
Entrez Gene	6401, 6403 (Human)		
SwissProt	P16581, P16109 (Human)		
Directions for Use	<ol style="list-style-type: none"> 1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube. 2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light. 3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells. 		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB

www.cohesionbio.com

ORDER

order@cohesionbio.com

SUPPORT

techsupport@cohesionbio.com

CUSTOM

custom@cohesionbio.com

Product Data Sheet

4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.
5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.
6. Add appropriate amount of fluorescent-labeled Streptavidin and incubate for 20 minutes away from light at room temperature.
7. Repeat step 5.
8. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB

www.cohesionbio.com

ORDER

order@cohesionbio.com

SUPPORT

techsupport@cohesionbio.com

CUSTOM

custom@cohesionbio.com