

## **Product Data Sheet**

# Anti-F4/80 Antibody-APC labled

Catalog #	Source	e Reactivity	Applications		
CFB8001	Rat	М	IF, FC		
Description		Rat monoclonal antibody Al	PC labled to F4/80		
Immunogen		Murine macrophages			
Purification		The antibody was purified b	y affinity chromatography.		
Specificity		Recognizes mouse F4/80			
Clonality		Monoclonal (clone: BM8)			
Conjugation		APC			
Form		Rat IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.			
Dilution					
Gene Symbol		ADGRE1			
Alternative Names		EMR1; TM7LN3; EGF-like module-containing mucin-like hormone receptor-like 1;			
		EGF-like module receptor 1	EMR1 hormone receptor; Adhesion G protein-coupled		
		receptor E1; F4/80			
Entrez Gene		13733 (Mouse)			
SwissProt		Q61549 (Mouse)			
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 µl labeled antiboo	ly to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	tes at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	0 rpm centrifugation for 5 minutes, discard the		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

错误!未指定文件名。

错误!未指定文件名。

错误!未指定文件名。

错误!未指定文件名。

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com