

Product Data Sheet

Anti-CD10 Antibody-APC labled

Catalog #	Source	Reactivity	Applications			
CFB8484	Mouse	H H	IF, FC			
Description	Description Mouse monoclonal antibody APC labled to CD10					
Immunogen		Native purified human CD10.				
Purification		The antibody was purified by affinity chromatography.				
Specificity		Recognizes human CD10				
Clonality		Monoclonal (clone: CB-CALLA)				
Conjugation		APC				
Form		Mouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Dilution		10 μl / assay				
Gene Symbol		MME				
Alternative Names		EPN; Neprilysin; Atriopeptidase; Common acute lymphocytic leukemia antigen;				
		CALLA; Enkephalinase; Neutral endopeptidase 24.11; NEP; Neutral endopeptidase;				
		Skin fibroblast elastase; SF	; CD antigen CD10			
Entrez Gene		4311 (Human)				
SwissProt		P08473 (Human)				
Directions for Use		1. Take 100 μ l peripheral bl	ood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.				
		2. Add 10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole			
		blood, incubate for 20 minu	ites at room temperature away from light.			
		3. Add 2 ml RBC lysis buffer	, incubate for 10 minutes away from light after mixing,			
		dissolve red blood cells.				
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com