

Product Data Sheet

Anti-CD11a Antibody-APC labled

| Catalog # | Source | e Reactivity | Applications | |
|--------------------|--------|--|---|--|
| CFB8488 | Mouse | - | IF, FC | |
| Description | | Mouse monoclonal antibod | | |
| Immunogen | | Native purified human CD12 | | |
| Purification | | The antibody was purified b | | |
| Specificity | | Recognizes human CD11a | | |
| Clonality | | Monoclonal (clone: Cris-3) | | |
| Conjugation | | APC | | |
| Form | | | in PBS, pH 7.3, and 0.02% sodium azide. | |
| | | | in PBS, pH 7.5, and 0.02% souldin azide. | |
| Dilution | | 10 μl / assay | | |
| Gene Symbol | | ITGAL | | |
| Alternative Names | | CD11A; Integrin alpha-L; CD11 antigen-like family member A; Leukocyte adhesion | | |
| | | glycoprotein LFA-1 alpha ch | ain; LFA-1A; Leukocyte function-associated molecule 1 | |
| | | alpha chain; CD antigen CD1 | 1a | |
| Entrez Gene | | 3683 (Human) | | |
| SwissProt | | P20701 (Human) | | |
| Directions for Use | | 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5 | | |
| | | ml tube. | | |
| | | 2. Add 10 μl labeled antiboo | y to the bottom of flow tube mixing with the whole | |
| | | blood, incubate for 20 minu | tes at room temperature away from light. | |
| | | 3. Add 2 ml RBC lysis buffer, | incubate for 10 minutes away from light after mixing, | |
| | | dissolve red blood cells. | | |
| | | | 0 rpm centrifugation for 5 minutes, discard the | |
| | | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com