

## **Product Data Sheet**

# Anti-CD73 Antibody-APC labled

Catalog #	Source	Reactivity	Applications		
CFB8552	Mouse	Н	IF, FC		
Description		Mouse monoclonal antiboo	dy APC labled to CD73		
Immunogen		Native purified human CD7	73.		
Purification		The antibody was purified	by affinity chromatography.		
Specificity		Recognizes human CD73			
Clonality		Monoclonal (clone: AD2)			
Conjugation		APC			
Form		Mouse IgG1. Liquid in PBS,	pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution		10 µl / assay			
Gene Symbol		NT5E			
Alternative Names		NT5; NTE; 5'-nucleotidase; 5'-NT; Ecto-5'-nucleotidase; CD antigen CD73			
Entrez Gene		4907 (Human)			
SwissProt		P21589 (Human)			
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 µl labeled antibo	dy to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 min	utes at room temperature away from light.		
		3. Add 2 ml RBC lysis buffe	r, incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 10	00 rpm centrifugation for 5 minutes, discard the		
		supernatant.			
		5. Add 2 ml PBS wash buffe	er to resuspend the cells, then 1000 rpm centrifugation		
Application kow		M/D Mastern blat III Insperior	histochomistry IF Immunofluoroscopeo FC Flow outomatry IC		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com