

Product Data Sheet

Anti-CD81 Antibody-APC labled

| Catalog # | Source | Reactivity | Applications | | |
|--------------------|--------|--|--|--|--|
| CFB8557 | Mouse | Н | IF, FC | | |
| Description | Ν | Nouse monoclonal antiboo | dy APC labled to CD81 | | |
| Immunogen | Ν | lative purified human CD8 | 1. | | |
| Purification | Т | he antibody was purified | by affinity chromatography. | | |
| Specificity | R | Recognizes human CD81 | | | |
| Clonality | Ν | Aonoclonal (clone: 1.3.3.2 | 2) | | |
| Conjugation | A | APC | | | |
| Form | | Mouse IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide. | | | |
| Dilution | 1 | .0 μl / assay | | | |
| Gene Symbol | C | CD81 | | | |
| Alternative Names | | TAPA1; TSPAN28; CD81 antigen; 26 kDa cell surface protein TAPA-1; Target of the | | | |
| | а | ntiproliferative antibody 1 | ; Tetraspanin-28; Tspan-28; CD antigen CD81 | | |
| Entrez Gene | | 975 (Human) | | | |
| SwissProt | | P60033 (Human) | | | |
| Directions for Use | | 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5 | | | |
| | n | nl tube. | | | |
| | 2 | . Add 10 μl labeled antibo | dy to the bottom of flow tube mixing with the whole | | |
| | b | lood, incubate for 20 min | utes at room temperature away from light. | | |
| | 3 | . Add 2 ml RBC lysis buffer | r, incubate for 10 minutes away from light after mixing, | | |
| | d | lissolve red blood cells. | | | |
| | 4 | . Sample tube is set to 10 | 00 rpm centrifugation for 5 minutes, discard the | | |
| | S | upernatant. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com