

# **Product Data Sheet**

## Anti-CD71 Antibody-APC labled

Catalog #	Source	Reactivity	Applications			
CFB8604	Mouse	Н	IF, FC			
Description		Mouse monoclonal antibo	dy APC labled to CD71			
Immunogen		NALM-6 pre-B cell line				
Purification		The antibody was purified by affinity chromatography.				
Specificity		Recognizes human CD71				
Clonality		Monoclonal (clone: RVS-10	))			
Conjugation		APC				
Form		Mouse IgG1 kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Dilution		10 μl / assay				
Gene Symbol		TFRC				
Alternative Names		Transferrin receptor protein 1; TR; TfR; TfR1; Trfr; T9; p90; CD71				
Entrez Gene		7037 (Human)				
SwissProt		P02786 (Human)				
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
		ml tube.				
		2. Add 10 μl labeled antibc	ody to the bottom of flow tube mixing with the whole			
		blood, incubate for 20 min	utes at room temperature away from light.			
		3. Add 2 ml RBC lysis buffe	r, incubate for 10 minutes away from light after mixing,			
		dissolve red blood cells.				
		4. Sample tube is set to 10	00 rpm centrifugation for 5 minutes, discard the			
		supernatant.				
		5. Add 2 ml PBS wash buffe	er to resuspend the cells, then 1000 rpm centrifugation			
Application kow		M/P Mostorn blot III Immune	shistashamistry IE Immunaflyarassansa EC Elaw sytamatry IC			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com