

## **Product Data Sheet**

# Anti-CD40 Antibody-APC labled

Catalog #	Source	Poactivity	Applications	
	Source		Applications	
CFB8796	Rat	Μ	IF, FC	
Description		Armenian Hamster monoc	lonal antibody APC labled to CD40	
Immunogen		(BALB/c x NZB)F1 mouse-c	lerived lymphoma WEHI 231	
Purification		The antibody was purified	by affinity chromatography.	
Specificity		Recognizes mouse CD40		
Clonality		Monoclonal (clone: 1C10)		
Conjugation		APC		
Form		Rat IgG2a kappa. Liquid in	PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.	
Dilution		10 μl / assay		
Gene Symbol		CD40		
Alternative Names		TNFRSF5; Tumor necrosis factor receptor superfamily member 5; B-cell surface		
		antigen CD40; Bp50; CD40	L receptor; CDw40; CD40	
Entrez Gene		21939 (Mouse)		
SwissProt		P27512 (Mouse)		
<b>Directions for Use</b>		1. Take 100 µl peripheral b	lood anticoagulated by EDTA and add to the bottom of 5	
		ml tube.		
		2. Add 10 µl labeled antibo	ody to the bottom of flow tube mixing with the whole	
		blood, incubate for 20 min	utes at room temperature away from light.	
		3. Add 2 ml RBC lysis buffe	r, incubate for 10 minutes away from light after mixing,	
		dissolve red blood cells.		
		4. Sample tube is set to 10	00 rpm centrifugation for 5 minutes, discard the	
		supernatant.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com