

# **Product Data Sheet**

# Anti-CD71 Antibody-APC/Cy7 labled

Catalog #	Source	Reactivity	Applications		
CFD8604	Mouse	e H	IF, FC		
Description		Mouse monoclonal antiboo	ly APC/Cy7 labled to CD71		
Immunogen		NALM-6 pre-B cell line			
Purification		The antibody was purified b	by affinity chromatography.		
Specificity		Recognizes human CD71			
Clonality		Monoclonal (clone: CY1G4)			
Conjugation		APC/Cy7			
Form		Mouse IgG2a kappa. Liquid	in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution		10 µl / assay			
Gene Symbol		TFRC			
Alternative Names		Transferrin receptor protein 1; TR; TfR; TfR1; Trfr; T9; p90; CD71			
Entrez Gene		7037 (Human)			
SwissProt		P02786 (Human)			
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 $\mu l$ labeled antibo	dy to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	ites at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer	; incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the		
		supernatant.			
		5. Add 2 ml PBS wash buffe	r to resuspend the cells, then 1000 rpm centrifugation		
Application kov		M/D Mastern blat III Insurance	histochemistry IF Immunofluorosconce FC Flow externetry IC		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com