

# **Product Data Sheet**

## Rat IgG2b Isotype Control-FITC labled

Source	e Reactivity Applications			
Rat	IF, FC			
	Rat IgG2b Isotype Control-FITC labled can be used as an isotype control in flow			
	cytometry and other applications.			
	The antibody was purified by affinity chromatography.			
	Monoclonal (clone: LTF-2)			
	FITC			
	Rat IgG2b. Liquid in PBS, pH 7.3, and 0.02% sodium azide.			
	10 μl / assay			
mes				
Jse	1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
	ml tube.			
	2. Add 10 $\mu l$ labeled antibody to the bottom of flow tube mixing with the whole			
	blood, incubate for 20 minutes at room temperature away from light.			
	3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing,			
	dissolve red blood cells.			
	4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the			
	supernatant.			
	Rat			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com