

## **Product Data Sheet**

# Anti-CD40 Antibody-FITC labled

Catalog #	Source	Reactivity	Applications		
CFE8527	Mouse	н	IF, FC		
Description		Mouse monoclonal antibod	y FITC labled to CD40		
Immunogen		Native purified human CD4	).		
Purification		The antibody was purified b	y affinity chromatography.		
Specificity		Recognizes human CD40			
Clonality		Monoclonal (clone: 3A8)			
Conjugation		FITC			
Form		Mouse IgG2b. Liquid in PBS	pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution		10 μl / assay			
Gene Symbol		CD40			
Alternative Names		TNFRSF5; Tumor necrosis factor receptor superfamily member 5; B-cell surface			
		antigen CD40; Bp50; CD40L	receptor; CDw40; CD antigen CD40		
Entrez Gene		958 (Human)			
SwissProt		P25942 (Human)			
Directions for	<sup>.</sup> Use	1. Take 100 $\mu$ l peripheral blo	ood anticoagulated by EDTA and add to the bottom of 5		
		ml tube.			
		2. Add 10 $\mu$ l labeled antibod	ly to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	tes at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	0 rpm centrifugation for 5 minutes, discard the		
		supernatant.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com