

# **Product Data Sheet**

# Anti-CD103 Antibody-FITC labled

| Catalog #                 | Source | Reactivity   | Applications  |  |  |
|---------------------------|--------|--|---|--|--|
| CFE8621                   | Mouse  | Н  | IF, FC  |  |  |
| Description               |        | Mouse monoclonal antiboo   | ly FITC labled to CD103                                 |  |  |
| Immunogen                 |        | HTLV-1 induced human T co  | ell line MAPS16   |  |  |
| Purification              |        | The antibody was purified  | by affinity chromatography.                             |  |  |
| Specificity               |        | Recognizes human CD103   |   |  |  |
| Clonality                 |        | Monoclonal (clone: Ber-AC  | ۲8)   |  |  |
| Conjugation               |        | FITC   |   |  |  |
| Form                      |        | Mouse IgG1 kappa. Liquid i   | n PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.        |  |  |
| Dilution                  |        | 10 µl / assay  |   |  |  |
| Gene Symbol               |        | ITGAE  |   |  |  |
| Alternative Names         |        | Integrin alpha-E; HML-1 antigen; Integrin alpha-IEL; Mucosal lymphocyte 1 antigen;     |   |  |  |
|                           |        | CD103  |   |  |  |
| Entrez Gene               |        | 3682 (Human)   |   |  |  |
| SwissProt                 |        | P38570 (Human)   |   |  |  |
| <b>Directions for Use</b> |        | 1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5 |   |  |  |
|                           |        | ml tube.   |   |  |  |
|                           |        | 2. Add 10 $\mu$ l labeled antibo   | dy to the bottom of flow tube mixing with the whole     |  |  |
|                           |        | blood, incubate for 20 min   | ites at room temperature away from light.               |  |  |
|                           |        | 3. Add 2 ml RBC lysis buffer   | , incubate for 10 minutes away from light after mixing, |  |  |
|                           |        | dissolve red blood cells.  |   |  |  |
|                           |        | 4. Sample tube is set to 100   | 00 rpm centrifugation for 5 minutes, discard the        |  |  |
|                           |        | supernatant.   |   |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com