

# **Product Data Sheet**

# Anti-CD161 Antibody-FITC labled

Catalog #	Source	e Reactivity	Applications	
CFE8658	Mouse	e H	IF, FC	
Description		Mouse monoclonal antibod	y FITC labled to CD161	
Immunogen		Human NK cells		
Purification		The antibody was purified b	y affinity chromatography.	
Specificity		Recognizes human CD161		
Clonality		Monoclonal (clone: HP-3G1	))	
Conjugation		FITC		
Form		Mouse IgG1 kappa. Liquid ir	PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.	
Dilution		10 µl / assay		
Gene Symbol		KLRB1		
Alternative Names		CLEC5B; NKRP1A; Killer cell lectin-like receptor subfamily B member 1; C-type lectin		
		domain family 5 member B;	HNKR-P1a; NKR-P1A; Natural killer cell surface protein	
		P1A; CD161		
Entrez Gene		3820 (Human)		
SwissProt		Q12918 (Human)		
<b>Directions for Use</b>		1. Take 100 μl peripheral blo	od anticoagulated by EDTA and add to the bottom of 5	
		ml tube.		
		2. Add 10 $\mu$ l labeled antibod	y to the bottom of flow tube mixing with the whole	
		blood, incubate for 20 minu	tes at room temperature away from light.	
		3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,	
		dissolve red blood cells.		
		4. Sample tube is set to 100	O rpm centrifugation for 5 minutes, discard the	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com