

Product Data Sheet

Anti-MPO Antibody-FITC labled

| Catalog # | Source | Reactivity | Applications | | |
|--------------------|--------------|--|--|--|--|
| CFE8763 I | Mouse | Н | IF, FC | | |
| Description | M | Mouse monoclonal antibody FITC labled to MPO | | | |
| Immunogen | | | | | |
| Purification | Th | The antibody was purified by affinity chromatography. | | | |
| Specificity | Re | Recognizes human MPO | | | |
| Clonality | M | Monoclonal (clone: MPO455-8E6) | | | |
| Conjugation | | | | | |
| Form | M | Mouse IgG1 kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide. | | | |
| Dilution | 10 | μl / assay | | | |
| Gene Symbol | М | PO | | | |
| Alternative Names | | Myeloperoxidase; MPO | | | |
| Entrez Gene | 43 | 4353 (Human) | | | |
| SwissProt | PO | P05164 (Human) | | | |
| Directions for U | se 1. | Take 100 μl peripheral bl | lood anticoagulated by EDTA and add to the bottom of 5 | | |
| | ml | tube. | | | |
| | 2. | Add 10 µl labeled antibo | dy to the bottom of flow tube mixing with the whole | | |
| | blo | ood, incubate for 20 min | utes at room temperature away from light. | | |
| | 3. | Add 2 ml RBC lysis buffe | r, incubate for 10 minutes away from light after mixing, | | |
| | dis | ssolve red blood cells. | | | |
| | 4. | Sample tube is set to 10 | 00 rpm centrifugation for 5 minutes, discard the | | |
| | su | pernatant. | | | |
| | 5. | Add 2 ml PBS wash buffe | er to resuspend the cells, then 1000 rpm centrifugation | | |
| Application key: F | - FLISA \//F | 8- Western blot IH- Immuno | histochemistry IE- Immunofluorescence EC- Flow cytometry IC- | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com