

Product Data Sheet

Anti-CD24 Antibody-FITC labled

| | Catalog # | Source | e Reactivity | Applications | | |
|---|--------------------|----------|--|--|--|--|
| | CFE8783 | Rat | Μ | IF, FC | | |
| | Description | | Rat monoclonal antibody FITC labled to CD24 | | | |
| | | | C57BL/10 mouse splenic T cells and concanavalin A-activated splenocytes | | | |
| | | | The antibody was purified by affinity chromatography. | | | |
| | | | Recognizes mouse CD24 | | | |
| Clonality Monoclonal (clone: 91) | | | | | | |
| | Conjugation | | FITC | | | |
| Form Dilution | | | Rat IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide. | | | |
| | | | 10 μl / assay | | | |
| | Gene Symbol | | CD24 | | | |
| | Alternative Na | mes | CD24A; Signal transducer C | CD24; Small cell lung carcinoma cluster 4 antigen; CD24 | | |
| Entrez Gene12484 (Mouse)SwissProtP24807 (Mouse) | | | 12484 (Mouse) | | | |
| | | | | | | |
| | Directions for | Use | 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5 | | | |
| | | ml tube. | | | | |
| | | | 2. Add 10 μl labeled antibody to the bottom of flow tube mixing with the whole | | | |
| | | | blood, incubate for 20 minutes at room temperature away from light. | | | |
| | | | 3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, | | | |
| | | | dissolve red blood cells. | | | |
| | | | 4. Sample tube is set to 100 | 00 rpm centrifugation for 5 minutes, discard the | | |
| | | | supernatant. | | | |
| | | | 5. Add 2 ml PBS wash buffe | er to resuspend the cells, then 1000 rpm centrifugation | | |
| | A sublication base | | | bisto show istra JE. Increase of Leave and a SC. Flow and a start to the | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com