

Product Data Sheet

Anti-CD161c Antibody-FITC labled

Catalog #	Source	Reactivity	Applications		
CFE8958	Mouse	М	IF, FC		
Description		Mouse monoclonal antibody	FITC labled to CD161c		
Immunogen					
Purification		The antibody was purified b	v affinity chromatography.		
Specificity		Recognizes mouse CD161c			
Clonality		Monoclonal (clone: PK136)			
Conjugation		FITC			
Form		Mouse IgG2a kappa. Liquid i	n PBS, pH 7.3, and 0.02% sodium azide.		
Dilution		10 μl / assay			
Gene Symbol		KLRB1C			
Alternative Names		Ly55c; Nkrp1c; Killer cell lectin-like receptor subfamily B member 1C; CD161			
		antigen-like family member	C; Lymphocyte antigen 55c; Ly-55c; NK1.1; NKR-P1.9;		
		NKR-P1C; Natural killer cell s	urface protein P1-40; NKR-P1 40; CD161c		
Entrez Gene		17059 (Mouse)			
SwissProt		P27814 (Mouse)			
Directions for	Use	1. Take 100 μ l peripheral blo	od anticoagulated by EDTA and add to the bottom of 5		
		ml tube.			
		2. Add 10 μ l labeled antibod	y to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minut	es at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 1000) rpm centrifugation for 5 minutes, discard the		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com