

Product Data Sheet

Anti-CD62E/CD62P Antibody-FITC labled

Catalog # Source Reactivity Applications

CFE8971 Mouse H IF, FC

Description Mouse monoclonal antibody FITC labled to CD62E/CD62P

Immunogen TNF stimulated human endothelial cells

Purification The antibody was purified by affinity chromatography.

Specificity Recognizes human CD62E/CD62P

Clonality Monoclonal (clone: 1.2B6)

Conjugation FITC

Form Mouse IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

Dilution 10 μl / assay

Gene Symbol SELE; SELP

Alternative Names ELAM1; E-selectin; CD62 antigen-like family member E; Endothelial leukocyte

adhesion molecule 1; ELAM-1; Leukocyte-endothelial cell adhesion molecule 2;

LECAM2; CD62E; GMRP; GRMP; P-selectin; CD62 antigen-like family member P;

Granule membrane protein 140; GMP-140; Leukocyte-endothelial cell adhesion

molecule 3; LECAM3; Platelet activation dependent granule-external membrane

protein; PADGEM; CD62P

Entrez Gene 6401, 6403 (Human)

SwissProt P16581, P16109 (Human)

Directions for Use 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5

ml tube.

2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole

blood, incubate for 20 minutes at room temperature away from light.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com





Product Data Sheet

- 3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells.
- 4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.
- 5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.
- 6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED