

**Anti-HLA-B27 Antibody-FITC labeled**

Catalog #	Source	Reactivity	Applications
CFE8974	Mouse	H	IF, FC
<b>Description</b>	Mouse monoclonal antibody FITC labeled to HLA-B27		
<b>Immunogen</b>	Immune complex precipitated from HLA-B27 positive cell line (Bordin)		
<b>Purification</b>	The antibody was purified by affinity chromatography.		
<b>Specificity</b>	Recognizes human HLA-B27		
<b>Clonality</b>	Monoclonal (clone: HLA-B27)		
<b>Conjugation</b>	FITC		
<b>Form</b>	Mouse IgG2a. Liquid in PBS, pH 7.3, and 0.02% sodium azide.		
<b>Dilution</b>	10 µl / assay		
<b>Gene Symbol</b>	HLA-B27		
<b>Alternative Names</b>	HLA-B27 protein		
<b>Entrez Gene</b>			
<b>SwissProt</b>	O19193 (Human)		
<b>Directions for Use</b>	<ol style="list-style-type: none"><li>1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube.</li><li>2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light.</li><li>3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells.</li><li>4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.</li><li>5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation</li></ol>		
<b>Application key:</b>	E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference		
<b>Species reactivity key:</b>	H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish		

**COHESION BIOSCIENCES LIMITED****WEB**[www.cohesionbio.com](http://www.cohesionbio.com)**ORDER**[order@cohesionbio.com](mailto:order@cohesionbio.com)**SUPPORT**[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)**CUSTOM**[custom@cohesionbio.com](mailto:custom@cohesionbio.com)

# Product Data Sheet

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

## Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

## COHESION BIOSCIENCES LIMITED

### WEB

[www.cohesionbio.com](http://www.cohesionbio.com)

### ORDER

[order@cohesionbio.com](mailto:order@cohesionbio.com)

### SUPPORT

[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

### CUSTOM

[custom@cohesionbio.com](mailto:custom@cohesionbio.com)