

Anti-Ly-49C/F/H/I Antibody-FITC labeled

Catalog #	Source	Reactivity	Applications
CFE8993	Syrian Hamster	M	IF, FC
Description	Syrian Hamster monoclonal antibody FITC labeled to Ly-49C/F/H/I		
Immunogen	IL-2-activated killer cells (LAK) from C57BL/6 mice		
Purification	The antibody was purified by affinity chromatography.		
Specificity	Recognizes mouse Ly-49C/F/H/I		
Clonality	Monoclonal (clone: 14B11)		
Conjugation	FITC		
Form	Syrian Hamster IgG. Liquid in PBS, pH 7.3, and 0.02% sodium azide.		
Dilution	10 µl / assay		
Gene Symbol	Klra3; Klra6; Klra8; Klra9		
Alternative Names	Ly-49c; Ly49C; Killer cell lectin-like receptor 3; 5E6; Lymphocyte antigen 49c; Ly-49c; Nk2.1; T-cell surface glycoprotein Ly-49C; Ly-49f; Ly49-f; Ly49F; Killer cell lectin-like receptor 6; Lymphocyte antigen 49f; Ly-49f; T-cell surface glycoprotein Ly-49F; Ly-49h; Ly49-h; Ly49H; Killer cell lectin-like receptor 8; Lymphocyte antigen 49h; Ly-49h; T-cell surface glycoprotein Ly-49H; Killer cell lectin-like receptor subfamily A member 9; Killer cell lectin-like receptor subfamily A member 9; Natural killer cell receptor Ly49C		
Entrez Gene	16634, 16637, 16639, 16640 (Mouse)		
SwissProt	Q64329, Q60653, Q60682, Q2TJJ8 (Mouse)		
Directions for Use	1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com

Product Data Sheet

2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light.
3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells.
4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.
5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.
6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB

www.cohesionbio.com

ORDER

order@cohesionbio.com

SUPPORT

techsupport@cohesionbio.com

CUSTOM

custom@cohesionbio.com