

Product Data Sheet

Anti-CD83 Antibody-PE labled

| Catalog # | Source | Reactivity | Applications | | |
|--------------------|--------|----------------------------------------------------------------------------------------|---------------------------------------------------------|--|--|
| CFF8558 | Mouse | Н | IF, FC | | |
| Description | | Mouse monoclonal antiboo | ly PE labled to CD83 | | |
| Immunogen | | Native purified human CD8 | 3. | | |
| Purification | | The antibody was purified | by affinity chromatography. | | |
| Specificity | | Recognizes human CD83 | | | |
| Clonality | | Monoclonal (clone: HB15e) | | | |
| Conjugation | | PE | | | |
| Form | | Mouse IgG1 kappa. Liquid i | n PBS, pH 7.3, and 0.02% sodium azide. | | |
| Dilution | | 10 µl / assay | | | |
| Gene Symbol | 1 | CD83 | | | |
| Alternative Names | | CD83 antigen; hCD83; B-cell activation protein; Cell surface protein HB15; CD | | | |
| | | antigen CD83 | | | |
| Entrez Gene | | 9308 (Human) | | | |
| SwissProt | | Q01151 (Human) | | | |
| Directions for Use | | 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5 | | | |
| | | ml tube. | | | |
| | | 2. Add 10 μl labeled antibo | dy to the bottom of flow tube mixing with the whole | | |
| | | blood, incubate for 20 min | utes at room temperature away from light. | | |
| | | 3. Add 2 ml RBC lysis buffer | ; incubate for 10 minutes away from light after mixing, | | |
| | | dissolve red blood cells. | | | |
| | | 4. Sample tube is set to 100 | 00 rpm centrifugation for 5 minutes, discard the | | |
| | : | supernatant. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com