

# **Product Data Sheet**

## Anti-CD25 Antibody-PE labled

Catalog #	Source	e Reactivity	Applications		
-					
CFF8784	Rat	Μ	IF, FC		
Description		Rat monoclonal antibody PE labled to CD25			
Immunogen		IL-2-dependent BALB/c mou	se helper T-cell clone HT-2		
Purification		The antibody was purified b	/ affinity chromatography.		
Specificity		Recognizes mouse CD25			
Clonality		Monoclonal (clone: PC61.5)			
Conjugation		PE			
Form		Rat IgG1 lambda. Liquid in P	BS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution		10 μl / assay			
Gene Symbol		IL2RA			
Alternative Names		Interleukin-2 receptor subunit alpha; IL-2 receptor subunit alpha; IL-2-RA; IL-2R			
		subunit alpha; IL2-RA; TAC a	ntigen; p55; CD25		
Entrez Gene		16184 (Mouse)			
SwissProt		P01590 (Mouse)			
<b>Directions for Use</b>		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 $\mu$ l labeled antibod	y to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	es at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	) rpm centrifugation for 5 minutes, discard the		
		supernatant.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com