

## Anti-CD69 Antibody-PE labeled

Catalog #	Source	Reactivity	Applications
CFF8824	Armenian Hamster	M	IF, FC
<b>Description</b>	Armenian Hamster monoclonal antibody PE labeled to CD69		
<b>Immunogen</b>	Mouse dendritic epidermal T cell line Y245		
<b>Purification</b>	The antibody was purified by affinity chromatography.		
<b>Specificity</b>	Recognizes mouse CD69		
<b>Clonality</b>	Monoclonal (clone: H1.2F3)		
<b>Conjugation</b>	PE		
<b>Form</b>	Armenian Hamster IgG. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
<b>Dilution</b>	10 µl / assay		
<b>Gene Symbol</b>	CD69		
<b>Alternative Names</b>	CLEC2C; Early activation antigen CD69; Activation inducer molecule; AIM; BL-AC/P26; C-type lectin domain family 2 member C; EA1; Early T-cell activation antigen p60; GP32/28; Leukocyte surface antigen Leu-23; MLR-3; CD69		
<b>Entrez Gene</b>	12515 (Mouse)		
<b>SwissProt</b>	P37217 (Mouse)		
<b>Directions for Use</b>	<ol style="list-style-type: none"> <li>1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube.</li> <li>2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light.</li> <li>3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells.</li> </ol>		

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

**WEB**  
[www.cohesionbio.com](http://www.cohesionbio.com)

**ORDER**  
[order@cohesionbio.com](mailto:order@cohesionbio.com)

**SUPPORT**  
[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

**CUSTOM**  
[custom@cohesionbio.com](mailto:custom@cohesionbio.com)

## Product Data Sheet

4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.
5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.
6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

### Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

#### WEB

[www.cohesionbio.com](http://www.cohesionbio.com)

#### ORDER

[order@cohesionbio.com](mailto:order@cohesionbio.com)

#### SUPPORT

[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

#### CUSTOM

[custom@cohesionbio.com](mailto:custom@cohesionbio.com)