

# **Product Data Sheet**

## **Anti-TDT Antibody-PE labled**

Catalog #	Source	Reactivity	Applications			
CFF8980	Mouse	н	IF, FC			
Description Mouse monoclonal antibody PE labled to TDT						
Immunogen		Recombinant full length hu	man TdT			
Purification		The antibody was purified b	by affinity chromatography.			
Specificity		Recognizes human TDT				
Clonality		Monoclonal (clone: 41A)				
Conjugation		PE				
Form		Mouse IgG2a kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.				
Dilution		10 μl / assay				
Gene Symbol		DNTT				
Alternative Names		TDT; DNA nucleotidylexotransferase; Terminal addition enzyme; Terminal				
		deoxynucleotidyltransferas	e; Terminal transferase			
Entrez Gene		1791 (Human)				
SwissProt		P04053 (Human)				
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
		ml tube.				
		2. Add 10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole			
		blood, incubate for 20 minu	utes at room temperature away from light.			
		3. Add 2 ml RBC lysis buffer	; incubate for 10 minutes away from light after mixing,			
		dissolve red blood cells.				
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the			
		supernatant.				

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com