

Product Data Sheet

Anti-TCR beta Antibody-PE/Cy5 labled

Catalog #	Source	Reactivity	Applications			
CFG8998	Armenian	М	IF, FC			
Hamster						
Description	Armen	iian hamster monocl	onal antibody PE/Cy5 labled to TCR beta			
Immunogen	Affinity	Affinity purified TCR from mouse DO-11.10 cells				
Purification	The an	The antibody was purified by affinity chromatography.				
Specificity	Recog	Recognizes mouse TCR beta				
Clonality	Monoo	Monoclonal (clone: H57-597)				
Conjugation	PE/Cy5	PE/Cy5				
Form	Armen	Armenian Hamster IgG. Liquid in PBS, pH 7.3, and 0.02% sodium azide.				
Dilution	10 µl /	10 μl / assay				
Gene Symbol	TCRB	TCRB				
Alternative Na	ames Tib; TC	Tib; TCRbeta				
Entrez Gene	21577	21577 (Mouse)				
SwissProt	(Mous	(Mouse)				
Directions for	Use 1. Take	1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5				
	ml tub	e.				
	2. Add	10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole			
	blood,	incubate for 20 min	utes at room temperature away from light.			
	3. Add	2 ml RBC lysis buffe	; incubate for 10 minutes away from light after mixing,			
	dissolv	e red blood cells.				
	4. Sam	ple tube is set to 10	00 rpm centrifugation for 5 minutes, discard the			
	superr	natant.				
	2. Add blood, 3. Add dissolv 4. Sam	10 μl labeled antibo incubate for 20 min 2 ml RBC lysis buffer ve red blood cells. aple tube is set to 100	utes at room temperature away from light.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com