

Product Data Sheet

Anti-CD4 Antibody-PerCP labled

| Catalog # | Source | e Reactivity | Applications | | |
|--------------------|--------|--|---|--|--|
| CFJ8525 | Mouse | e H | IF, FC | | |
| Description | | Mouse monoclonal antibo | dy PerCP labled to CD4 | | |
| Immunogen | | Native purified human CD4 | 1. | | |
| Purification | | The antibody was purified | by affinity chromatography. | | |
| Specificity | | Recognizes human CD4 | | | |
| Clonality | | Monoclonal (clone: Riv7) | | | |
| Conjugation | | PerCP | | | |
| Form | | Mouse IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide. | | | |
| Dilution | | 10 μl / assay | | | |
| Gene Symbol | | CD4 | | | |
| Alternative Names | | T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3; CD antigen CD4 | | | |
| Entrez Gene | | 920 (Human) | | | |
| SwissProt | | P01730 (Human) | | | |
| Directions for Use | | 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5 | | | |
| | | ml tube. | | | |
| | | 2. Add 10 μ l labeled antibo | ody to the bottom of flow tube mixing with the whole | | |
| | | blood, incubate for 20 min | utes at room temperature away from light. | | |
| | | 3. Add 2 ml RBC lysis buffe | r, incubate for 10 minutes away from light after mixing, | | |
| | | dissolve red blood cells. | | | |
| | | 4. Sample tube is set to 10 | 00 rpm centrifugation for 5 minutes, discard the | | |
| | | supernatant. | | | |
| | | 5. Add 2 ml PBS wash buffe | er to resuspend the cells, then 1000 rpm centrifugation | | |
| Application kow | | M/D Mastern blat III Immun | histochomistry IE Immunofluorosconso EC Elow automatry IC | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com