

## **Product Data Sheet**

# Anti-CD64 Antibody-PerCP/Cy5.5 labled

Catalog #	Source	Reactivity	Applications		
CFK8549	Mouse		IF, FC		
Description			dy PerCP/Cy5.5 labled to CD64		
Immunogen		Native purified human CD6			
Purification			by affinity chromatography.		
		Recognizes human CD64	by animity thromatography.		
Specificity		C			
Clonality		Monoclonal (clone: 10.1)			
Conjugation		PerCP/Cy5.5			
Form		Mouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.			
Dilution		10 µl / assay			
Gene Symbol		FCGR1A			
Alternative Names		FCG1; FCGR1; IGFR1; High affinity immunoglobulin gamma Fc receptor I; IgG Fc			
		receptor I; Fc-gamma RI; Fo	RI; Fc-gamma RIA; FcgammaRIa; CD antigen CD64		
Entrez Gene		2209 (Human)			
SwissProt		P12314 (Human)			
<b>Directions for Use</b>		1. Take 100 μl peripheral b	lood anticoagulated by EDTA and add to the bottom of 5		
		ml tube.			
		2. Add 10 µl labeled antibo	dy to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 min	utes at room temperature away from light.		
		3. Add 2 ml RBC lysis buffe	r, incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 10	00 rpm centrifugation for 5 minutes, discard the		
		supernatant.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com