

# **Product Data Sheet**

# Anti-CD79a Antibody-PerCP/Cy5.5 labled

Catalog # Sour	rce Reactivity	Applications			
CFK8606 Mou	use H	IF, FC			
Description	Mouse monoclonal antibody PerCP/Cy5.5 labled to CD79a				
Immunogen					
Purification	The antibody was purified by affinity chromatography.				
Specificity	Recognizes human CD79a				
Clonality	Monoclonal (clone: HM47)				
Conjugation	PerCP/Cy5.5				
Form	Mouse IgG1 kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Dilution	10 μl / assay	10 μl / assay			
Gene Symbol	CD79A				
Alternative Names	IGA; MB1; B-cell antigen receptor complex-associated protein alpha chain; Ig-alpha;				
	MB-1 membrane glycopr	otein; Membrane-bound immunoglobulin-associated			
	protein; Surface IgM-asso	ociated protein; CD79a			
Entrez Gene	973 (Human)				
SwissProt	P11912 (Human)				
Directions for Use	$1.$ Take 100 $\mu$ l peripheral blood anticoagulated by EDTA and add to the bottom of 5				
	ml tube.				
	2. Add 10 µl labeled antil	oody to the bottom of flow tube mixing with the whole			
	blood, incubate for 20 m	nutes at room temperature away from light.			
	3. Add 2 ml RBC lysis buf	er, incubate for 10 minutes away from light after mixing,			
	dissolve red blood cells.				
	4. Sample tube is set to 1	000 rpm centrifugation for 5 minutes, discard the			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com