

# **Product Data Sheet**

## Anti-CD107a Antibody-AF647 labled

Catalog #	Source	Reactivity	Applications		
CFN8623	Mouse	н	IF, FC		
Description Mouse monoclonal antibody AF647 labled to CD107a					
Immunogen		Human adult adherent per	pheral blood cells		
Purification		The antibody was purified by affinity chromatography.			
Specificity		Recognizes human CD107a			
Clonality		Monoclonal (clone: H4A3)			
Conjugation		AF647			
Form		Mouse IgG1 kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.			
Dilution		10 μl / assay			
Gene Symbol		LAMP1			
Alternative Names		Lysosome-associated membrane glycoprotein 1; LAMP-1; Lysosome-associated			
		membrane protein 1; CD10	7 antigen-like family member A; CD107a		
Entrez Gene		3916 (Human)			
SwissProt		P11279 (Human)			
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 $\mu$ l labeled antibo	dy to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	ites at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer	; incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the		
		supernatant.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com