

Product Data Sheet

Anti-CD11c Antibody-Pacific Blue labled

Catalog #	Source	Reactivity	Applications			
CFP8776	Armenian	Μ	IF, FC			
Hamster						
Description	Arme	nian Hamster monocl	onal antibody Pacific Blue labled to CD11c			
Immunogen	Mous	Mouse spleen dendritic cells				
Purification	The a	The antibody was purified by affinity chromatography.				
Specificity	Recog	Recognizes mouse CD11c				
Clonality	Mono	Monoclonal (clone: N418)				
Conjugation	Pacifi	Pacific Blue				
Form	Arme	Armenian Hamster IgG. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Dilution	10 µl	10 μl / assay				
Gene Symbol	ITGAX	ITGAX				
Alternative N	ames CD11	CD11C; Integrin alpha-X; CD11 antigen-like family member C; Leu M5; Leukocyte				
	adhes	sion glycoprotein p15	0,95 alpha chain; Leukocyte adhesion receptor p150,95;			
	CD11	с				
Entrez Gene	16412	16411 (Mouse)				
SwissProt	Q9QX	Q9QXH4 (Mouse)				
Directions for	Use 1. Tak	1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5				
	ml tu	be.				
	2. Ad	d 10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole			
	blood	l, incubate for 20 min	utes at room temperature away from light.			
	3. Ad	d 2 ml RBC lysis buffe	r, incubate for 10 minutes away from light after mixing,			
	dissol	ve red blood cells.				

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com