

# **Product Data Sheet**

## Anti-CD102 Antibody-Pacific Blue labled

Catalog #	Source	e Reactivity	Applications		
CFP8841	Rat	Μ	IF, FC		
Description		Rat monoclonal antibody P	acific Blue labled to CD102		
Immunogen		COS cells transfected with r	nouse ICAM-2 cDNA		
Purification		The antibody was purified b	by affinity chromatography.		
Specificity		Recognizes mouse CD102			
Clonality		Monoclonal (clone: 3C4)			
Conjugation		Pacific Blue			
Form		Rat IgG2a kappa. Liquid in I	PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution		10 μl / assay			
Gene Symbol		ICAM2			
Alternative Names		Intercellular adhesion molecule 2; ICAM-2; CD102			
Entrez Gene		15896 (Mouse)			
SwissProt		P35330 (Mouse)			
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 µl labeled antibo	dy to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	utes at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer	; incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the		
		supernatant.			
		5. Add 2 ml PBS wash buffe	r to resuspend the cells, then 1000 rpm centrifugation		
Application kov		M/D Mastern blat III Insperies	histochemistry IE Immunofluorosconco EC Elouy outomotry IC		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com