

Product Data Sheet

Anti-Ly-6C Antibody-Pacific Blue labled

| Catalog # | Source | e Reactivity | Applications | | |
|--------------------|--|--|--|--|--|
| CFP8995 | Rat | Μ | IF, FC | | |
| Description | n Rat monoclonal antibody Pacific Blue labled to Ly-6C | | | | |
| Immunogen | | L3 cloned CTL cells | | | |
| Purification | | The antibody was purified b | y affinity chromatography. | | |
| Specificity | | Recognizes mouse Ly-6C | | | |
| Clonality | | Monoclonal (clone: HK1.4) | | | |
| Conjugation | | Pacific Blue | | | |
| Form | | Rat IgG2c kappa. Liquid in P | BS, pH 7.3, and 0.02% sodium azide. | | |
| Dilution | | 10 μl / assay | | | |
| Gene Symbol | | Ly6c1 | | | |
| Alternative Names | | Lymphocyte antigen 6C1; Ly-6C1 | | | |
| Entrez Gene | | 17067 (Mouse) | | | |
| SwissProt | | P0CW02 (Mouse) | | | |
| Directions for Use | | 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5 | | | |
| | | ml tube. | | | |
| | | 2. Add 10 μl labeled antiboc | ly to the bottom of flow tube mixing with the whole | | |
| | | blood, incubate for 20 minu | tes at room temperature away from light. | | |
| | | 3. Add 2 ml RBC lysis buffer, | incubate for 10 minutes away from light after mixing, | | |
| | | dissolve red blood cells. | | | |
| | | 4. Sample tube is set to 100 | 0 rpm centrifugation for 5 minutes, discard the | | |
| | | supernatant. | | | |
| | | 5. Add 2 ml PBS wash buffer | to resuspend the cells, then 1000 rpm centrifugation | | |
| Application key | | W/D Western blot III Immunek | istochomistry IF, Immunofluorosconco, FC, Flow externatry IC | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com