

# **Product Data Sheet**

## Anti-TCR beta Antibody-Pacific Blue labled

Catalog #	Source	Reactivity	Applications			
CFP8998	Armenian	Μ	IF, FC			
	Hamster					
Description	Armen	Armenian hamster monoclonal antibody Pacific Blue labled to TCR beta				
Immunogen	Affinity	Affinity purified TCR from mouse DO-11.10 cells				
Purification	The an	The antibody was purified by affinity chromatography.				
Specificity	Recogr	Recognizes mouse TCR beta				
Clonality	Monoc	Monoclonal (clone: H57-597)				
Conjugation	Pacific	Pacific Blue				
Form	Armen	Armenian Hamster IgG. Liquid in PBS, pH 7.3, and 0.02% sodium azide.				
Dilution	10 µl /	10 μl / assay				
Gene Symbol	TCRB	TCRB				
Alternative Na	imes Tib; TC	Tib; TCRbeta				
Entrez Gene	21577	21577 (Mouse)				
SwissProt	(Mouse	(Mouse)				
Directions for	Use 1. Take	1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
	ml tub	e.				
	2. Add	10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole			
	blood,	incubate for 20 minu	ites at room temperature away from light.			
	3. Add	2 ml RBC lysis buffer	, incubate for 10 minutes away from light after mixing,			
	dissolv	e red blood cells.				
	4. Sam	ple tube is set to 100	00 rpm centrifugation for 5 minutes, discard the			
	supern	atant.				

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com