

Product Data Sheet

hsa-miR-208a-3p miRNA Antagomir

Catalog #	Source	Reactiv	vity Applications		
CIJ0050	Synthetic	c H			
Description	Sy	Synthetic miRNA Antagomir is used to inhibit the activity of target hsa-miR-208a-3p			
	r	nRNA.			
Specificity	А	ntagomir is ch	emically-modified single-strand miRNA inhibitor functioning by		
	b	locking miRNA	regulation of target gene expression efficiently. They are synthesized		
	to	o reduce the a	pility of endogenous miRNAs to silence target mRNA transcripts.		
	TI	hey can down-	regulate the corresponding endogenous miRNAs. Our miRNA		
	a	ntagomir is sin	gle-strand miRNA inhibitor carrying the chemically modifications		
	fı	unctioning by b	blocking miRNA regulation of target gene expression efficiently.		
Form	Ly	yophilized pow	/der		
Gene Symbol	h	sa-miR-208a-3	ρ		
Accession No.	. N	1IMAT0000241			
Components	T	This synthetic miRNA is based on the mature miRNA sequence. The strand of the			
	a	ntagomir has 2	2 phosphorothioates at the 5' end, 4 phosphorothioates, 1		
	cł	holesterol grou	up at the 3' end, and full-length nucleotide 2'-methoxy modification.		
	St	tability of miRI	NA antagomir appears to be significantly higher than miRNA		
	in	nhibitors. It e	whibits enhanced cellular uptake, stability and regulatory activity and		
	is	recommende	d for miRNA functional studies in vitro and in vivo.		
Directions for Use Briefly centrifuge tubes containing miRNA antagomir to e		e tubes containing miRNA antagomir to ensure that the miRNA			
	р	ellet is located	at the bottom of the tube. Dissolve miRNA antagomir to a		
	CC	onvenient stoc	k concentration using the recommended volume of DEPC H2O (or		
	R	Nase-free wat	er). For example: dissolve 10 nmol miRNA antagomir to 20 μM using		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com