

Product Data Sheet

hsa-miR-133a-3p miRNA Antagomir

Catalog #	Source	Reac	tivity Applications			
CIJ0100	Syntheti	c H				
Description	S	Synthetic miRNA Antagomir is used to inhibit the activity of target hsa-miR-133a-3p				
	n	nRNA.				
Specificity	А	Antagomir is chemically-modified single-strand miRNA inhibitor functioning by				
	b	locking miRN	NA regulation of target gene expression efficiently. They are synthesized			
	to	o reduce the	ability of endogenous miRNAs to silence target mRNA transcripts.			
	Т	hey can dow	n-regulate the corresponding endogenous miRNAs. Our miRNA			
	а	intagomir is s	ingle-strand miRNA inhibitor carrying the chemically modifications			
	fu	unctioning by	y blocking miRNA regulation of target gene expression efficiently.			
Form	L	yophilized po	owder			
Gene Symbol	h	ısa-miR-133a	-3p			
Accession No.	. N	ЛІМАТ00004	27			
Components	Т	This synthetic miRNA is based on the mature miRNA sequence. The strand of the				
	а	intagomir has	s 2 phosphorothioates at the 5' end, 4 phosphorothioates, 1			
	C	holesterol gr	oup at the 3' end, and full-length nucleotide 2'-methoxy modification.			
	S	tability of mi	RNA antagomir appears to be significantly higher than miRNA			
	ir	nhibitors. II	t exhibits enhanced cellular uptake, stability and regulatory activity and			
	is	s recomment	ded for miRNA functional studies in vitro and in vivo.			
Directions for	Directions for Use Briefly centrifuge tubes containing miRNA antagomir to ensure that t		uge tubes containing miRNA antagomir to ensure that the miRNA			
	р	ellet is locate	ed at the bottom of the tube. Dissolve miRNA antagomir to a			
	C	onvenient st	ock concentration using the recommended volume of DEPC H2O (or			
	R	Nase-free wa	ater). For example: dissolve 10 nmol miRNA antagomir to 20 μM using			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com