

## **Product Data Sheet**

# hsa-miR-520c-3p miRNA Antagomir

| Catalog #      | Source        | Reactivity         | Applications   |
|----------------|---------------|--------------------|--|
| CIJ0277        | Synthetic     | Н                  |  |
| Description    | Sy            | nthetic miRNA A    | ntagomir is used to inhibit the activity of target hsa-miR-520c-3p   |
|                | ml            | RNA.               |  |
| Specificity    | Ar            | tagomir is chem    | ically-modified single-strand miRNA inhibitor functioning by         |
|                | blo           | ocking miRNA rea   | gulation of target gene expression efficiently. They are synthesized |
|                | to            | reduce the abilit  | ry of endogenous miRNAs to silence target mRNA transcripts.          |
|                | Th            | ey can down-reg    | ulate the corresponding endogenous miRNAs. Our miRNA                 |
|                | an            | tagomir is single  | -strand miRNA inhibitor carrying the chemically modifications        |
|                | fu            | nctioning by bloc  | king miRNA regulation of target gene expression efficiently.         |
| Form           | Lyo           | ophilized powde    | r  |
| Gene Symbol    | hs            | a-miR-520c-3p      |  |
| Accession No.  | . М           | MAT0002846         |  |
| Components     | Th            | is synthetic miRI  | NA is based on the mature miRNA sequence. The strand of the          |
|                | an            | tagomir has 2 pł   | nosphorothioates at the 5' end, 4 phosphorothioates, 1               |
|                | ch            | olesterol group a  | at the 3' end, and full-length nucleotide 2'-methoxy modification.   |
|                | Sta           | ability of miRNA   | antagomir appears to be significantly higher than miRNA              |
|                | inl           | nibitors. It exhi  | bits enhanced cellular uptake, stability and regulatory activity and |
|                | is            | recommended fo     | or miRNA functional studies in vitro and in vivo.                    |
| Directions for | <b>Use</b> Br | efly centrifuge t  | ubes containing miRNA antagomir to ensure that the miRNA             |
|                | pe            | llet is located at | the bottom of the tube. Dissolve miRNA antagomir to a                |
|                | со            | nvenient stock co  | oncentration using the recommended volume of DEPC H2O (or            |
|                | RN            | lase-free water).  | For example: dissolve 10 nmol miRNA antagomir to 20 $\mu M$ using    |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com