

# **Product Data Sheet**

## hsa-miR-337-5p miRNA Antagomir

Catalog #	Source	Reactivity	Applications
CIJ0583	Synthetic	: H	
Description	Sy	nthetic miRNA A	ntagomir is used to inhibit the activity of target hsa-miR-337-5p
	m	RNA.	
Specificity	А	ntagomir is chem	ically-modified single-strand miRNA inhibitor functioning by
	bl	ocking miRNA reg	gulation of target gene expression efficiently. They are synthesized
	to	reduce the abilit	y of endogenous miRNAs to silence target mRNA transcripts.
	TI	ney can down-reg	ulate the corresponding endogenous miRNAs. Our miRNA
	aı	ntagomir is single	-strand miRNA inhibitor carrying the chemically modifications
	fu	inctioning by bloc	king miRNA regulation of target gene expression efficiently.
Form	Ly	ophilized powder	
Gene Symbol	h	sa-miR-337-5p	
Accession No.	. N	IIMAT0004695	
Components	TI	nis synthetic miRN	IA is based on the mature miRNA sequence. The strand of the
	aı	ntagomir has 2 ph	osphorothioates at the 5' end, 4 phosphorothioates, 1
	cł	nolesterol group a	t the 3' end, and full-length nucleotide 2'-methoxy modification.
	St	ability of miRNA	antagomir appears to be significantly higher than miRNA
	in	hibitors. It exhi	bits enhanced cellular uptake, stability and regulatory activity and
	is	recommended for	or miRNA functional studies in vitro and in vivo.
<b>Directions for</b>	<b>Directions for Use</b> Briefly centrifuge tubes containing miRNA antagomir to ensure that the mi		
	р	ellet is located at	the bottom of the tube. Dissolve miRNA antagomir to a
	CC	onvenient stock co	oncentration using the recommended volume of DEPC H2O (or
	R	Nase-free water).	For example: dissolve 10 nmol miRNA antagomir to 20 $\mu M$ using

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

500 μl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com