

## **Product Data Sheet**

## hsa-miR-4704-3p miRNA Antagomir

| Catalog #     | Source   | Re  | activity   | Applications   |  |
|---------------|--|---|--|--|--|
| CIJ1537       | Synthet  | tic H   |  |  |  |
| Description   | n Synth  |   | thetic miRNA Antagomir is used to inhibit the activity of target hsa-miR-4704-3p |  |  |
|               |  | mRNA.   |  |  |  |
| Specificity   |  | Antagomir is chemically-modified single-strand miRNA inhibitor functioning by |  |  |  |
|               |  | blocking mi   | iRNA regulatio   | on of target gene expression efficiently. They are synthesized |  |
|               |  | to reduce the ability of endogenous miRNAs to silence target mRNA transcript  |  |  |  |
|               | ,  | They can down-regulate the corresponding endogenous miRNAs. Our miRNA         |  |  |  |
|               |  | antagomir i   | is single-stran  | d miRNA inhibitor carrying the chemically modifications        |  |
|               |  | functioning   | by blocking r  | niRNA regulation of target gene expression efficiently.        |  |
| Form          |  | Lyophilized   | powder   |  |  |
| Gene Symbol   |  | hsa-miR-47  | 04-3p  |  |  |
| Accession No. |  | MIMAT001  | 9804   |  |  |
| Components    | ,  | This synthe   | tic miRNA is b   | pased on the mature miRNA sequence. The strand of the          |  |
|               |  | antagomir l   | has 2 phospho  | prothioates at the 5' end, 4 phosphorothioates, 1              |  |
|               |  | cholesterol   | group at the   | 3' end, and full-length nucleotide 2'-methoxy modification.    |  |
|               | Stability of miRNA antagomir appears to be significantly higher than i |   | omir appears to be significantly higher than miRNA                               |  |  |
|               |  | inhibitors.   | It exhibits e  | nhanced cellular uptake, stability and regulatory activity and |  |
|               |  | is recomme  | ended for miR  | NA functional studies in vitro and in vivo.                    |  |

**Directions for Use** 

Briefly centrifuge tubes containing miRNA antagomir to ensure that the miRNA pellet is located at the bottom of the tube. Dissolve miRNA antagomir to a convenient stock concentration using the recommended volume of DEPC H2O (or RNase-free water). For example: dissolve 10 nmol miRNA antagomir to 20  $\mu$ M using

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**





## **Product Data Sheet**

500 µl DEPC H2O (or RNase-free water). Pipette the solution up and down 3-5 times (or vortex briefly). Briefly centrifuge tubes containing miRNA antagomir to ensure that the solution is collected at the bottom of the tube. Aliquot the miRNA antagomir into small volumes and store at ≤ -20°C. miRNA antagomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

**COHESION BIOSCIENCES LIMITED**