

Product Data Sheet

Recombinant Anti-hnRNP U Rabbit mAb

Catalog # Source Reactivity Applications

CMA4332 Rabbit H WB, IF/IC

Description Recombinant rabbit monoclonal antibody to hnRNP U

Immunogen KLH-conjugated synthetic peptide encompassing a sequence within human hnRNP

U. The exact sequence is proprietary.

Purification The antibody was purified by immunogen affinity chromatography.

Specificity Recognizes endogenous levels of hnRNP U protein

Clonality Monoclonal

Conjugation

Form Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.

Dilution WB (1/500 - 1/1000), IF/IC (1/50 - 1/200)

Gene Symbol HNRNPU

Alternative Names HNRPU; SAFA; U21.1; Heterogeneous nuclear ribonucleoprotein U; hnRNP U;

Scaffold attachment factor A; SAF-A; p120; pp120

Entrez Gene 3192 (Human)

SwissProt Q00839 (Human)

Storage/Stability Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid

freeze/thaw cycles.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference


Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED


WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com



Product Data Sheet



Western blot analysis of hnRNP U expression in Hela (A), THP1 (B), HepG2 (C) whole cell lysates. (Predicted band size: 90 kD; Observed band size: 120 kD)



Immunofluorescent analysis of hnRNP U staining in A375 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish