

Product Data Sheet

rno-miR-497-5p miRNA Agomir

| Catalog # | Source | Reactivity | Applications | | | |
|------------------|------------------|--|--|--|--|--|
| CMS0231 | Synthetic | R | | | | |
| Description | Synth | etic miRNA Agomir is | used to regulate the expression of target rno-miR-497-5p | | | |
| | mRN | ۹. | | | | |
| Specificity | Agom | ir is chemically-modif | ied double-strand miRNA mimics which can mimic | | | |
| | matu | re endogenous miRNA | s after transfection into cells. They can up-regulate the | | | |
| | endo | genous miRNA activity | by utilizing the natural miRNA machinery. Our miRNA | | | |
| | agom | ir is designed to mimi | c mature miRNAs, and chemically modified to increase | | | |
| | their | stability and activity. | | | | |
| Form | Lyoph | nilized powder | | | | |
| Gene Symbol | rno-n | niR-497-5p | | | | |
| Accession No. | MIM | AT0003383 | | | | |
| Components | This s | This synthetic miRNA is based on the mature miRNA sequence. The antisense strand | | | | |
| | of the | e agomir has 2 phosph | orothioates at the 5' end, 4 phosphorothioates, 1 | | | |
| | chole | sterol group at the 3' | end, and full-length nucleotide 2'-methoxy modification. | | | |
| | lt exh | ibits enhanced cellula | r uptake, stability and regulatory activity in vivo. | | | |
| Directions for U | Jse Brief | y centrifuge tubes cor | taining miRNA agomir to ensure that the miRNA pellet is | | | |
| | locate | ed at the bottom of th | e tube. Dissolve miRNA agomir to a convenient stock | | | |
| | conce | entration using the rec | ommended volume of DEPC H2O (or RNase-free water). | | | |
| | For ex | kample: dissolve 5 nm | ol miRNA agomir to 20 μ M using 250 μ l DEPC H2O (or | | | |
| | RNase | e-free water). Pipette | the solution up and down 3-5 times (or vortex briefly). | | | |
| | Brief | y centrifuge tubes cor | taining miRNA agomir to ensure that the solution is | | | |
| | collec | ted at the bottom of t | he tube. Aliquot the miRNA agomir into small volumes | | | |
| Application key: | - FLISA \W/R_ \M | lestern blot IH- Immuno | histochemistry IE- Immunofluorescence EC- Flow cytometry IC- | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

and store at \leq -20°C. miRNA agomir is stable (for 6 months under the specified storage condition). For best results, use in 3 months and limit freeze-thaw events for each tube no more than five times.

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year. Avoid freeze-thaw cycles after reconstitution.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com