

Product Data Sheet

Anti-TRAP220 Antibody

Catalog #	Source	e Reactivity	Applications	
CPA1917	Rabbit	H, M, R, Mk	WB, IH, IF/IC	
Description		Rabbit polyclonal antibody to	TRAP220	
Immunogen		KLH-conjugated synthetic peptide encompassing a sequence within the center		
		region of human TRAP220. T	ne exact sequence is proprietary.	
Purification		The antibody was purified by	immunogen affinity chromatography.	
Specificity		Recognizes endogenous levels of TRAP220 protein.		
Clonality		Polyclonal		
Conjugation				
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
		and 0.01% sodium azide.		
Dilution		WB (1/500 - 1/1000), IH (1/100) - 1/200), IF/IC (1/100 - 1/500)	
Gene Symbol		MED1		
Alternative N	ames	ARC205; CRSP1; CRSP200; DF	IP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A;	
		TRAP220; TRIP2; Mediator of	RNA polymerase II transcription subunit 1;	
		Activator-recruited cofactor 2	05 kDa component; ARC205; Mediator complex	
		subunit 1; Peroxisome prolife	rator-activated receptor-binding protein; PBP;	
		PPAR-binding protein; Thyroi	d hormone receptor-associated protein complex 220	
		kDa component; Trap220; Th	yroid receptor-interacting protein 2; TR-interacting	
		protein 2; TRIP-2; Vitamin D	eceptor-interacting protein complex component	
		DRIP205; p53 regulatory prot	ein RB18A	
Entrez Gene		5469 (Human); 19014 (Mous	e)	
SwissProt		Q15648 (Human); Q925J9 (N	ouse)	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com




Product Data Sheet


Storage/Stability

Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid


freeze/thaw cycles.



Western blot analysis of TRAP220 expression in H446 (A) whole cell lysates. (Predicted band size: 168 kD; Observed band size: 220 kD)



Immunohistochemical analysis of TRAP220 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of TRAP220 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com